

On the vertices of the k -additive core

Michel GRABISCH^{a,*}, Pedro MIRANDA^b

^a*University of Paris I – Panthéon-Sorbonne, Centre d'Économie de la Sorbonne,
106-112 Bd de l'Hôpital, 75013 Paris, France*

^b*Universidad Complutense de Madrid, Plaza de Ciencias, 3, 28040 Madrid, Spain*

Abstract

The core of a game v on N , which is the set of additive games ϕ dominating v such that $\phi(N) = v(N)$, is a central notion in cooperative game theory, decision making and in combinatorics, where it is related to submodular functions, matroids and the greedy algorithm. In many cases however, the core is empty, and alternative solutions have to be found. We define the k -additive core by replacing additive games by k -additive games in the definition of the core, where k -additive games are those games whose Möbius transform vanishes for subsets of more than k elements. For a sufficiently high value of k , the k -additive core is nonempty, and is a convex closed polyhedron. Our aim is to establish results similar to the classical results of Shapley and Ichiishi on the core of convex games (corresponds to Edmonds' theorem for the greedy algorithm), which characterize the vertices of the core.

Key words: cooperative games; core; k -additive games; vertices

* Corresponding author. Tel (+33) 1-44-07-82-85, Fax (+33) 1-44-07-83-01.

Email addresses: `michel.grabisch@univ-paris1.fr` (Michel GRABISCH),
`pmiranda@mat.ucm.es` (Pedro MIRANDA).

1 Introduction

Given a finite set N of n elements, and a set function $v : 2^N \rightarrow \mathbb{R}$ vanishing on the empty set (called hereafter a *game*), its *core* $\mathcal{C}(v)$ is the set of additive set functions ϕ on N such that $\phi(S) \geq v(S)$ for every $S \subseteq N$, and $\phi(N) = v(N)$. Whenever nonempty, the core is a convex closed bounded polyhedron.

In many fields, the core is a central notion which has deserved a lot of studies. In cooperative game theory, it is the set of imputations for players so that no subcoalition has interest to form [18]. In decision making under uncertainty, where games are replaced by *capacities* (monotonic games), it is the set of probability measures which are coherent with the given representation of uncertainty [19]. More on a combinatorial point of view, cores of convex games are also known as base polytopes associated to supermodular functions [13,9], for which the greedy algorithm is known to be a fundamental optimization technique. Many studies have been done along this line, e.g., by Faigle and Kern for the matching games [8], and cost games [7]. In game theory, which will be our main framework here, related notions are the selectope [3], and the Shapley value with many of its variations on combinatorial structures (see, e.g., [1]).

It is a well known fact that the core is nonempty if and only if the game is balanced [4]. In the case of emptiness, an alternative solution has to be found. One possibility is to search for games more general than additive ones, for example k -additive games and capacities proposed by Grabisch [10]. In short, k -additive games have their Möbius transform vanishing for subsets of more than k elements, so that 1-additive games are just usual additive games. Since any game is a k -additive game for some k (possibly $k = n$), the k -additive core, i.e., the set of dominating k -additive games, is never empty provided k is high enough. The authors have justified this definition in the framework of cooperative game theory [15]. Briefly speaking, an element of

the k -additive core implicitly defines by its Möbius transform an imputation (possibly negative), which is now defined on groups of at most k players, and no more on individuals. By definition of the k -additive core, the total worth assigned to a coalition will be always greater or equal to the worth the coalition can achieve by itself; however, the precise sharing among players has still to be decided (e.g., by some bargaining process) among each group of at most k players.

In game theory, elements of the core are imputations for players, and thus it is natural that they fulfill monotonicity. We call monotonic core the core restricted to monotonic games (capacities). We will see in the sequel that the core is usually unbounded, while the monotonic core is not.

The properties of the (classical) core are well known, most remarkable being the result characterizing the core of convex games, where the set of vertices is exactly the set of additive games induced by maximal chains (or equivalently by permutations on N) in the Boolean lattice $(2^N, \subseteq)$. This has been shown by Shapley [17], and later Ichiishi proved the converse implication [12]. This result is also known in the field of matroids, since vertices of the base polytope can be found by a greedy algorithm.

A natural question arises: is it possible to generalize the Shapley-Ichiishi theorem for k -additive (monotonic) cores? More precisely, can we find the set of vertices for some special classes of games? Are they induced by some generalization of maximal chains? The paper shows that the answer is more complex than expected. It is possible to define notions similar to permutations and maximal chains, so as to generate vertices of the k -additive core of $(k + 1)$ -monotone games, a result which is a true generalization of the Shapley-Ichiishi theorem, but this does not permit to find all vertices of the core. A full analytical description of vertices seems to be difficult to find, but we completely explicit the case $k = n - 1$.

After a preliminary section introducing necessary concepts, Section 3 presents our basic ingredients, that is, orders on subsets of at most k elements, and achievable families, which play the role of maximal chains in the classical case. Then Section 4 presents the main result on the characterization of vertices for $(k + 1)$ -monotone games induced by achievable families.

2 Preliminaries

Throughout the paper, $N := \{1, \dots, n\}$ denotes a set of n elements (players in a game, nodes of a graph, etc.). We use indifferently 2^N or $\mathcal{P}(N)$ for denoting the set of subsets of N , and the set of subsets of N containing at most k elements is denoted by $\mathcal{P}^k(N)$, while $\mathcal{P}_*^k(N) := \mathcal{P}^k(N) \setminus \{\emptyset\}$. For convenience, subsets like $\{i\}, \{i, j\}, \{2\}, \{2, 3\}, \dots$ are written in the compact form $i, ij, 2, 23$ and so on.

A *game* on N is a function $v : 2^N \rightarrow \mathbb{R}$ such that $v(\emptyset) = 0$. The set of games on N is denoted by $\mathcal{G}(N)$. For any $A \in 2^N \setminus \{\emptyset\}$, the *unanimity game centered on A* is defined by $u_A(B) := 1$ iff $B \supseteq A$, and 0 otherwise.

A game v on N is said to be:

- (i) *additive* if $v(A \cup B) = v(A) + v(B)$ whenever $A \cap B = \emptyset$;
- (ii) *convex* if $v(A \cup B) + v(A \cap B) \geq v(A) + v(B)$, for all $A, B \subseteq N$;
- (iii) *monotone* if $v(A) \leq v(B)$ whenever $A \subseteq B$;
- (iv) *k -monotone* for $k \geq 2$ if for any family of k subsets $A_1, \dots, A_k$, it holds

$$v\left(\bigcup_{i=1}^k A_i\right) \geq \sum_{\substack{K \subseteq \{1, \dots, k\} \\ K \neq \emptyset}} (-1)^{|K|+1} v\left(\bigcap_{j \in K} A_j\right)$$

- (v) *infinitely monotone* if it is k -monotone for all $k \geq 2$.

Convexity corresponds to 2-monotonicity. Note that k -monotonicity implies

78 k' -monotonicity for all $2 \leq k' \leq k$. Also, $(n - 2)$ -monotone games on N
79 are infinitely monotone [2]. The set of monotone games on N is denoted by
80 $\mathcal{MG}(N)$, while the set of infinitely monotone games is denoted by $\mathcal{G}_\infty(N)$.

Let v be a game on N . The *Möbius transform* of v [16] (also called *dividends* of v , see Harsanyi [11]) is a function $m : 2^N \rightarrow \mathbb{R}$ defined by:

$$m(A) := \sum_{B \subseteq A} (-1)^{|A \setminus B|} v(B), \quad \forall A \subseteq N.$$

The Möbius transform is invertible since one can recover v from m by:

$$v(A) = \sum_{B \subseteq A} m(B), \quad \forall A \subseteq N.$$

81 If v is an additive game, then m is non null only for singletons, and $m(\{i\}) =$
82 $v(\{i\})$. The Möbius transform of u_A is given by $m(A) = 1$ and m is 0 otherwise.

83 A game v is said to be k -additive [10] for some integer $k \in \{1, \dots, n\}$ if
84 $m(A) = 0$ whenever $|A| > k$, and there exists some A such that $|A| = k$, and
85 $m(A) \neq 0$.

86 Clearly, 1-additive games are additive. The set of games on N being at most
87 k -additive (resp. infinitely monotone games at most k -additive) is denoted by
88 $\mathcal{G}^k(N)$ (resp. $\mathcal{G}_\infty^k(N)$). As above, replace $\mathcal{G}$ by $\mathcal{MG}$ if monotone games are
89 considered instead.

90 We recall the fundamental following result.

91 **Proposition 1** [5] *Let v be a game on N . For any $A, B \subseteq N$, with $A \subseteq B$,*
92 *we denote $[A, B] := \{L \subseteq N \mid A \subseteq L \subseteq B\}$.*

(i) *Monotonicity is equivalent to*

$$\sum_{L \in [i, B]} m(L) \geq 0, \quad \forall B \subseteq N, \quad \forall i \in B.$$

(ii) For $2 \leq k \leq n$, k -monotonicity is equivalent to

$$\sum_{L \in [A, B]} m(L) \geq 0, \quad \forall A, B \subseteq N, A \subseteq B, \quad 2 \leq |A| \leq k.$$

93 Clearly, a monotone and infinitely monotone game has a nonnegative Möbius
94 transform.

The *core* of a game v is defined by:

$$\mathcal{C}(v) := \{\phi \in \mathcal{G}^1(N) \mid \phi(A) \geq v(A), \quad \forall A \subseteq N, \text{ and } \phi(N) = v(N)\}.$$

95 A *maximal chain* in 2^N is a sequence of subsets $A_0 := \emptyset, A_1, \dots, A_{n-1}, A_n := N$
96 such that $A_i \subset A_{i+1}$, $i = 0, \dots, n-1$. The set of maximal chains of 2^N is
97 denoted by $\mathcal{M}(2^N)$.

To each maximal chain $C := \{\emptyset, A_1, \dots, A_n = N\}$ in $\mathcal{M}(2^N)$ corresponds
a unique permutation σ on N such that $A_1 = \sigma(1)$, $A_2 \setminus A_1 = \sigma(2)$, $\dots$,
 $A_n \setminus A_{n-1} = \sigma(n)$. The set of all permutations over N is denoted by $\mathfrak{S}(N)$.
Let v be a game. Each permutation σ (or maximal chain C) induces an additive
game ϕ^σ (or ϕ^C) on N defined by:

$$\phi^\sigma(\sigma(i)) := v(\{\sigma(1), \dots, \sigma(i)\}) - v(\{\sigma(1), \dots, \sigma(i-1)\})$$

or

$$\phi^C(\sigma(i)) := v(A_i) - v(A_{i-1}), \quad \forall i \in N.$$

98 with the above notation. The following is immediate.

Proposition 2 *Let v be a game on N , and C a maximal chain of 2^N . Then*

$$\phi^C(A) = v(A), \quad \forall A \in C.$$

99 **Theorem 1** *The following propositions are equivalent.*

100 (i) v is a convex game.

101 (ii) All additive games ϕ^σ , $\sigma \in \mathfrak{S}(N)$, belong to the core of v .

102 (iii) $\mathcal{C}(v) = \text{co}(\{\phi^\sigma\}_{\sigma \in \mathfrak{S}(N)})$.

103 (iv) $\text{ext}(\mathcal{C}(v)) = \{\phi^\sigma\}_{\sigma \in \mathfrak{S}(N)}$,

104 where $\text{co}(\cdot)$ and $\text{ext}(\cdot)$ denote respectively the convex hull of some set, and the

105 extreme points of some convex set.

106 (i) $\Rightarrow$ (ii) and (i) $\Rightarrow$ (iv) are due to Shapley [17], while (ii) $\Rightarrow$ (i) was proved

107 by Ichiishi [12].

A natural extension of the definition of the core is the following. For some integer $1 \leq k \leq n$, the k -additive core of a game v is defined by:

$$\mathcal{C}^k(v) := \{\phi \in \mathcal{G}^k(N) \mid \phi(A) \geq v(A), \quad \forall A \subseteq N, \phi(N) = v(N)\}.$$

In a context of game theory where elements of the core are imputations, it is natural to consider that monotonicity must hold, i.e., the imputation allocated to some coalition $A \in \mathcal{P}_*^k(N)$ is larger than for any subset of A . We call it the *monotone k -additive core*:

$$\mathcal{MC}^k(v) := \{\phi \in \mathcal{MG}^k(N) \mid \phi(A) \geq v(A), \quad \forall A \subseteq N, \phi(N) = v(N)\}.$$

We introduce also the *core of k -additive infinitely monotone games*:

$$\mathcal{C}_\infty^k(v) := \{\phi \in \mathcal{G}_\infty^k(N) \mid \phi(A) \geq v(A), \quad \forall A \subseteq N, \text{ and } \phi(N) = v(N)\}.$$

108 The latter is introduced just for mathematical convenience, and has no clear

109 application. Note that $\mathcal{C}(v) = \mathcal{C}^1(v) = \mathcal{C}_\infty^1(v)$.

110 3 Orders on $\mathcal{P}_*^k(N)$ and achievable families

111 As our aim is to give a generalization of the Shapley-Ichiishi results, we need

112 counterparts of permutations and maximal chains. These are given in this sec-

tion. Exact connections between our material and permutations and maximal chains will be explicated at the end of this section. First, we introduce total orders on subsets of at most k elements as a generalization of permutations.

We denote by $\prec$ a total (strict) order on $\mathcal{P}_*^k(N)$, $\preceq$ denoting the corresponding weak order.

(i) $\prec$ is said to be *compatible* if for all $A, B \in \mathcal{P}_*^k(N)$, $A \prec B$ if and only if $A \cup C \prec B \cup C$ for all $C \subseteq N$ such that $A \cup C, B \cup C \in \mathcal{P}_*^k(N)$, $A \cap C = B \cap C = \emptyset$.

(ii) $\prec$ is said to be $\subseteq$ -*compatible* if $A \subset B$ implies $A \prec B$.

(iii) $\prec$ is said to be *strongly compatible* if it is compatible and $\subseteq$ -compatible.

We introduce the *binary order* $\prec^2$ on 2^N as follows. To any subset $A \subseteq N$ we associate an integer $\eta(A)$, whose binary code is the indicator function of A , i.e., the i th bit of $\eta(A)$ is 1 if $i \in A$, and 0 otherwise. For example, with $n = 5$, $\{1, 3\}$ and $\{4\}$ have binary codes 00101 and 01000 respectively, hence $\eta(\{1, 3\}) = 5$ and $\eta(\{4\}) = 8$. Then $A \prec^2 B$ if $\eta(A) < \eta(B)$. This gives

$$\begin{aligned} 1 \prec^2 2 \prec^2 12 \prec^2 3 \prec^2 13 \prec^2 23 \prec^2 123 \prec^2 4 \prec^2 14 \prec^2 24 \prec^2 \\ 124 \prec^2 34 \prec^2 134 \prec^2 234 \prec^2 1234 \prec^2 5 \prec^2 \dots \end{aligned} \quad (1)$$

Note the recursive nature of $\prec^2$. Obviously, $\prec^2$ is a strongly compatible order, as well as all its restrictions to $\mathcal{P}_*^k(N)$, $k = 1, \dots, n-1$.

We introduce now a generalization of maximal chains associated to permutations. Let $\prec$ be a total order on $\mathcal{P}_*^k(N)$. For any $B \in \mathcal{P}_*^k(N)$, we define

$$\mathcal{A}(B) := \{A \subseteq N \mid [A \supseteq B] \text{ and } [\forall K \subseteq A \text{ s.t. } K \in \mathcal{P}_*^k(N), \text{ it holds } K \preceq B]\}$$

the *achievable family* of B .

EXAMPLE 1: Consider $n = 3$, $k = 2$, and the following order: $1 \prec 2 \prec 12 \prec$

13 $\prec$ 23 $\prec$ 3. Then

$$\begin{aligned}\mathcal{A}(1) &= \{1\}, & \mathcal{A}(2) &= \{2\}, & \mathcal{A}(12) &= \{12\}, & \mathcal{A}(13) &= \mathcal{A}(23) = \emptyset, \\ \mathcal{A}(3) &= \{3, 13, 23, 123\}.\end{aligned}$$

126 **Proposition 3** $\{\mathcal{A}(B)\}_{B \in \mathcal{P}_*^k(N)}$ is a partition of $\mathcal{P}(N) \setminus \{\emptyset\}$.

127 **Proof:** Let $\emptyset \neq C \in \mathcal{P}(N)$. It suffices to show that there is a unique $B \in$
 128 $\mathcal{P}_*^k(N)$ such that $C \in \mathcal{A}(B)$. Let $K_1, K_2, \dots, K_p$ be the nonempty collection
 129 of subsets of C in $\mathcal{P}_*^k(N)$, assuming $K_1 \prec K_2 \prec \dots \prec K_p$. Then $C \in \mathcal{A}(K_p)$
 130 is the unique possibility, since any B outside the collection will fail to satisfy
 131 the condition $B \subseteq C$, and any $B \neq K_p$ inside the collection will fail to satisfy
 132 the condition $K_p \preceq B$. ■

133

134 **Proposition 4** For any $B \in \mathcal{P}_*^k(N)$ such that $\mathcal{A}(B) \neq \emptyset$, $(\mathcal{A}(B), \subseteq)$ is an
 135 inf-semilattice, with bottom element B .

136 **Proof:** If $\mathcal{A}(B) \neq \emptyset$, any $C \in \mathcal{A}(B)$ contains B , hence every $K \subseteq B \subseteq C$,
 137 $K \in \mathcal{P}_*^k(N)$, is such that $K \preceq B$. Hence $B \in \mathcal{A}(B)$, and it is the smallest
 138 element.

139 Let $K, K' \in \mathcal{A}(B)$, assuming $\mathcal{A}(B)$ contains at least 2 elements (otherwise,
 140 we are done). $K \in \mathcal{A}(B)$ is equivalent to $K \supseteq B$ and any $L \subseteq K$, $L \in \mathcal{P}_*^k(N)$
 141 is such that $L \preceq B$. The same holds for K' . Therefore, $K \cap K' \supseteq B$, and if
 142 $L \subseteq K \cap K'$, $L \in \mathcal{P}_*^k(N)$, then $L \subseteq K$ and $L \subseteq K'$, which entails $L \preceq B$.
 143 Hence $K \cap K' \in \mathcal{A}(B)$. ■

144

145 From the above proposition we deduce:

146 **Corollary 1** Let $B \in \mathcal{P}_*^k(N)$ and $\prec$ be some total order on $\mathcal{P}_*^k(N)$. Then

147 $\mathcal{A}(B) \neq \emptyset$ if and only if for all $C \in \mathcal{P}_*^k(N)$, $C \subseteq B$ implies $C \preceq B$. Conse-
 148 quently, if $|B| = 1$ then $\mathcal{A}(B) \neq \emptyset$.

149 **Corollary 2** $\mathcal{A}(B) \neq \emptyset$ for all $B \in \mathcal{P}_*^k(N)$ if and only if $\prec$ is $\subseteq$ -compatible.

150 It is easy to build examples where achievable families are not lattices.

151 **EXAMPLE 2:** Consider $n = 4, k = 2$ and the following order: 2, 3, 24, 12,
 152 4, 13, 34, 1, 23, 14. Then $\mathcal{A}(23) = \{23, 123, 234\}$, and $1234 \notin \mathcal{A}(23)$ since
 153 $14 \succ 23$.

154 Assuming $\mathcal{A}(B)$ is a lattice, we denote by $\check{B}$ its top element.

155 **Proposition 5** Let $\prec$ be a total order on $\mathcal{P}_*^k(N)$. Consider $B \in \mathcal{P}_*^k(N)$ such
 156 that $\mathcal{A}(B)$ is a lattice. Then it is a Boolean lattice isomorphic to $(\mathcal{P}(\check{B} \setminus B), \subseteq)$.


157 **Proof:** It suffices to show that $\mathcal{A}(B) = \{B \cup K \mid K \subseteq \check{B} \setminus B\}$. Taking
 158 $\check{K} := \check{B} \setminus B$, we have $B \cup \check{K} \in \mathcal{A}(B)$. Hence, any $L \subseteq B \cup \check{K}$, $L \in \mathcal{P}_*^k(N)$, is
 159 such that $L \preceq B$. This is a fortiori true for $L \subseteq B \cup K$, $L \in \mathcal{P}_*^k(N)$, $\forall K \subseteq \check{K}$.
 160 Hence $B \cup K$ belongs to $\mathcal{A}(B)$, for all $K \subseteq \check{K}$. ■

161

162 **Proposition 6** Assume $\prec$ is compatible. For any $B \in \mathcal{P}_*^k(N)$ such that
 163 $\mathcal{A}(B) \neq \emptyset$, $\mathcal{A}(B)$ is the Boolean lattice $[B, \check{B}]$.

164 **Proof:** If $\mathcal{A}(B)$ is a lattice, we know by Prop. 5 that it is a Boolean lattice
 165 with bottom element B . Since we know that $\mathcal{A}(B)$ is an inf-semilattice by
 166 Prop. 4, it remains to show that $K, K' \in \mathcal{A}(B)$ implies $K \cup K' \in \mathcal{A}(B)$.
 167 Assume $K \cup K' \notin \mathcal{A}(B)$. Then there exists $L \subseteq K \cup K'$, $L \in \mathcal{P}_*^k(N)$ such that
 168 $L \succ B$. Necessarily, $L \setminus K \neq \emptyset$, otherwise $L \subseteq K$ and $K \in \mathcal{A}(B)$ imply $L \prec B$,
 169 a contradiction. Similarly, $L \setminus K' \neq \emptyset$. Moreover, $L \not\subseteq B$ since $\mathcal{A}(B) \neq \emptyset$ (see
 170 Cor. 1).

171 We consider $D := L \setminus K$, not empty by definition of L . Since $L \setminus D \subseteq K$ and
 172 $L \setminus D \in \mathcal{P}_*^k(N)$, we have $L \setminus D \preceq B$, with strict inequality since $L \setminus D$ has
 173 elements outside $K \cap K'$, hence outside B (see Figure below).



174
 175 Suppose first that $|B| < k$, and let $D := \{i, j, \dots\}$. We have $B \cup l \in \mathcal{P}_*^k(N)$
 176 and $B \cup l \subseteq K'$ for any $l \in D$, which implies $B \cup l \prec B$. Taking $l = i$, by
 177 compatibility, $L \setminus D \prec B$ implies $(L \setminus D) \cup i \prec B \cup i \prec B$. By compatibility
 178 again, $(L \setminus D) \cup i \prec B$ implies $(L \setminus D) \cup i \cup j \prec B \cup j \prec B$. Continuing the
 179 process till all elements of D have been taken, we finally end with $L \prec B$, a
 180 contradiction.

181 Secondly, assume that $|B| = k$. Take $K'' \subset B$ such that $K'' \supseteq L \cap B$ and
 182 $|K'' \cup D| = k$, which is always possible by assumption. Since $K'' \subset B \subseteq K$
 183 and $K'' \in \mathcal{P}_*^k(N)$, we have $K'' \prec B$. Then

- 184 (i) Either $L \setminus D \prec K'' \prec B$. By compatibility, $L \setminus D \prec K''$ implies $L \prec K'' \cup D$.
 185 Since $K'' \cup D \in \mathcal{P}_*^k(N)$ and $K'' \cup D \subseteq K'$, we deduce that $K'' \cup D \prec B$,
 186 hence $L \prec B$, a contradiction.
 187 (ii) Or $K'' \prec L \setminus D \prec B$. Since $(L \setminus D) \cap (B \setminus K'') = \emptyset$, from compatibility
 188 $K'' \prec L \setminus D$ implies $B = K'' \cup (B \setminus K'') \prec (L \setminus D) \cup (B \setminus K'')$. We have by
 189 assumption $|(L \setminus D) \cup (B \setminus K'')| = |L| \leq k$ and $(L \setminus D) \cup (B \setminus K'') \subseteq K$,
 190 from which we deduce $(L \setminus D) \cup (B \setminus K'') \prec B$. Hence we get $B \prec B$, a
 191 contradiction.

192 ■

193
 194 The following example shows that compatibility is not a necessary condition.

EXAMPLE 3: Consider $n = 4$, $k = 2$, and the following order: 1, 3, 2, 12, 23, 13, 4, 14, 24, 34. This order is not compatible since $3 \prec 2$ and $12 \prec 13$.

We obtain:

$$\begin{aligned} \mathcal{A}(1) &= 1, & \mathcal{A}(3) &= 3, & \mathcal{A}(2) &= 2, & \mathcal{A}(12) &= 12, & \mathcal{A}(23) &= 23, & \mathcal{A}(13) &= \{13, 123\}, \\ \mathcal{A}(4) &= 4, & \mathcal{A}(14) &= 14, & \mathcal{A}(24) &= \{24, 124\}, & \mathcal{A}(34) &= \{34, 134, 234, 1234\}. \end{aligned}$$

195 All families are lattices.

196 In the above example, $\prec$ was $\subseteq$ -compatible. However, this is not enough to
197 ensure that achievable families are lattices, as shown by the following example.

EXAMPLE 4: Let us consider the following $\subseteq$ -compatible order with $n = 4$ and $k = 2$:

$$3 \prec 4 \prec 34 \prec 2 \prec 24 \prec 1 \prec 13 \prec 12 \prec 23 \prec 14.$$

198 Then $\mathcal{A}(23) = \{23, 123, 234\}$.

199 We give some fundamental properties of achievable families when they are
200 lattices, in particular of their top elements.

201 **Proposition 7** *Assume $\prec$ is compatible, and consider a nonempty achiev-*
202 *able family $\mathcal{A}(B)$, with top element $\check{B}$. Then $\{\mathcal{A}(B_i) \mid B_i \in \mathcal{P}_*^k(N), B_i \subseteq$*
203 *$\check{B}, \mathcal{A}(B_i) \neq \emptyset\}$ is a partition of $\mathcal{P}(\check{B}) \setminus \{\emptyset\}$.*

204 **Proof:** We know by Prop. 3 that all $\mathcal{A}(B_i)$'s are disjoint. It remains to show
205 that (1) any $K \subseteq \check{B}$ is in some $\mathcal{A}(B_i)$, $B_i \subseteq \check{B}$, and (2) conversely that any
206 K in such $\mathcal{A}(B_i)$ is a subset of $\check{B}$.

207 (1) Assume $K \in \mathcal{A}(B_i)$, $B_i \not\subseteq \check{B}$. Then $B_i \subseteq K \subseteq \check{B}$, a contradiction.

208 (2) Assume $K \in \mathcal{A}(B_i)$, $B_i \subseteq \check{B}$, and $K \not\subseteq \check{B}$. Then there exists $l \in K$ such
209 that $l \notin \check{B}$ (and hence not in B_i). Note that this implies $B_i \cup l \prec B_i$, provided
210 $|B_i| < k$. First we show that $l \prec j$ for any $j \in B_i$. Since $K \supseteq B_i \cup \{l\}$, we

deduce that for any $j \in B_i$, $\{j, l\} \prec B_i$ and $l \prec B_i$. If $B_i = \{j\}$, we can further deduce that $l \prec j$. Otherwise, if $B_i = \{j, j'\}$, from $\{j, l\} \prec \{j, j'\}$ and $\{j', l\} \prec \{j, j'\}$, by compatibility $l \prec j$ and $l \prec j'$. Generalizing the above, we conclude that $l \prec j$ for all $j \in B_i$.

Next, if $l \notin \check{B}$, it means that for some $B' \subseteq \check{B}$ such that $B' \cup l \in \mathcal{P}_*^k(N)$, we have $B' \cup l \succ B$ (otherwise l should belong to $\check{B}$). We prove that $B' \not\supseteq B_i$. The case $|B_i| = k$ is obvious, let us consider $|B_i| < k$. Suppose on the contrary that $B' = B_i \cup L$, $L \subseteq N \setminus B_i$. Then $B_i \cup l \prec B_i$ implies that $B' \cup l = B_i \cup l \cup L \prec B_i \cup L \prec B$, the last inequality coming from $B_i \cup L \subseteq \check{B}$, $B_i \cup L \in \mathcal{P}_*^k(N)$. But $B' \cup l \succ B$, a contradiction.

Choose any $j \in B_i \setminus B'$. Since $j \succ l$, we deduce $B' \cup j \succ B' \cup l \succ B$, but since $B' \cup j \subseteq \check{B}$ and $B' \cup j \in \mathcal{P}_*^k(N)$, it follows that $B' \cup j \prec B$, a contradiction.

■

Proposition 8 *Let $\prec$ be a compatible order on $\mathcal{P}_*^k(N)$. For any $B \in \mathcal{P}_*^k(N)$ such that $\mathcal{A}(B)$ is nonempty, putting $\check{B} := \{i_1, \dots, i_l\}$ with $i_1 \prec \dots \prec i_l$, then necessarily there exists $j \in \{1, \dots, l\}$ such that $B = \{i_j, \dots, i_l\}$.*

Proof: Assume $\check{B} \neq B$, otherwise we have simply $j = 1$. Consider i_j the element in B with the lowest index in the list $\{1, \dots, l\}$. Let us prove that all successors $i_{j+1}, \dots, i_l$ are also in B . Assume $j < l$ (otherwise we are done), and suppose that $i_{j'} \notin B$ for some $j < j' \leq l$. Then by compatibility, $B = (B \setminus i_j) \cup i_j \prec (B \setminus i_j) \cup i_{j'}$. Since $(B \setminus i_j) \cup i_{j'} \subseteq \check{B}$ and $(B \setminus i_j) \cup i_{j'} \in \mathcal{P}_*^k(N)$, the converse inequality should hold. ■

Proposition 9 *Assume that $\prec$ is strongly compatible. Then for all $B \subseteq N$, $|B| < k$, $\check{B} = B$.*

237 **Proof:** By Prop. 6 and Cor. 2, we know that $\mathcal{A}(B)$ is a Boolean lattice with
 238 top element denoted by $\check{B}$. Suppose that $\check{B} \neq B$. Then there exists $i \in \check{B} \setminus B$,
 239 and $B \cup i \in \mathcal{A}(B)$. Remark that $|B \cup i| \leq k$ and $\mathcal{A}(B \cup i) \ni B \cup i$ by Prop. 6
 240 and Cor. 2 again. This contradicts the fact that the achievable families form
 241 a partition of $\mathcal{P}_*^k(N)$ (Prop. 3). ■

242

Proposition 10 *Let $\prec$ be a strongly compatible order on $\mathcal{P}_*^k(N)$, and assume
 w.l.o.g. that $1 \prec 2 \prec \dots \prec n$. Then the collection $\check{\mathcal{B}}$ of $\check{B}$'s is given by:*

$$\check{\mathcal{B}} = \left\{ \{1, 2, \dots, l\} \cup \{j_1, \dots, j_{k-1}\} \mid l = 1, \dots, n - k + 1 \right. \\ \left. \text{and } \{j_1, \dots, j_{k-1}\} \subseteq \{l + 1, \dots, n\} \right\} \cup \left\{ A \subseteq N \mid |A| < k \right\}.$$

243 *If $\prec$ is compatible, then $\check{\mathcal{B}}$ is a subcollection of the above, where some subsets
 244 of at most $k - 1$ elements may be absent.*

245 **Proof:** From Prop. 9, we know that $\check{\mathcal{B}}$ contains all subsets having less than
 246 k elements. This proves the right part of “ $\cup$ ” in $\check{\mathcal{B}}$. By Prop. 9 again, the left
 247 part uniquely comes from those B 's of exactly k elements. Take such a B . From
 248 Prop. 8, we know that $\check{B}$ cannot contain elements ranked after the last one
 249 of B in the sequence $1, 2, \dots, n$. In other words, letting $B := \{l, j_1, \dots, j_{k-1}\}$,
 250 with l the lowest ranked element, we know that $\check{B} = B' \cup \{l, j_1, \dots, j_{k-1}\}$,
 251 with all elements of B' ranked before l . It remains to show that necessarily
 252 B' contains all elements from 1 to l excluded. Assume $j \notin B'$, $1 \leq j < l$.
 253 Then it should exist $K \in \mathcal{P}_*^k(N)$, $j \in K \subseteq \check{B} \cup j$, such that $K \succ B$. Since
 254 $|B| = k$, it cannot be that $K \supseteq B$, so that say $j' \in B$ is not in K . Hence
 255 we have $j \prec j'$, and by compatibility, $K = (K \setminus j) \cup j \prec (K \setminus j) \cup j'$. Now,
 256 $(K \setminus j) \cup j' \in \mathcal{P}_*^k(N)$ and $(K \setminus j) \cup j' \subseteq \check{B}$, which entails $(K \setminus j) \cup j' \prec B$, so
 257 that $K \prec B$, a contradiction.

258 Finally, consider that $\prec$ is only compatible. Then by Cor. 2, there exists $B \in$

259 $\mathcal{P}_*^k(N)$ such that $\mathcal{A}(B) = \emptyset$. This implies that there exist some proper subsets
 260 of B in $\mathcal{P}_*^k(N)$ ranked after B , let us call K the last ranked such subset. Then
 261 $|K| < k$, and $\mathcal{A}(K) \neq \{K\}$ since it contains at least B , because all subsets of
 262 B are ranked before K by definition of K . Hence K does not belong to $\check{\mathcal{B}}$. ■
 263

We finish this section by explaining why achievable families induced by orders on $\mathcal{P}_*^k(N)$ are generalizations of maximal chains induced by permutations. Taking $k = 1$, $\mathcal{P}_*^1(N) = N$, and total orders on singletons coincide with permutations on N . Trivially, any order on N is strongly compatible, so that all achievable families are nonempty lattices. Denoting by σ the permutation corresponding to $\prec$, i.e., $\sigma(1) \prec \sigma(2) \prec \dots \prec \sigma(n)$, then

$$\mathcal{A}(\{\sigma(j)\}) = [\{\sigma(j)\}, \{\sigma(1), \dots, \sigma(j)\}],$$

264 i.e., the top element $\{\sigma(j)\}$ is $\{\sigma(1), \dots, \sigma(j)\}$. Then the collection of all top
 265 elements $\{\sigma(j)\}$ is exactly the maximal chain associated to σ .

266 4 Vertices of $\mathcal{C}^k(v)$ induced by achievable families

267 Let us consider a game v and its k -additive core $\mathcal{C}^k(v)$. We suppose hereafter
 268 that $\mathcal{C}^k(v) \neq \emptyset$, which is always true for a sufficiently high k . Indeed, taking
 269 at worst $k = n$, $v \in \mathcal{C}^n(v)$ always holds.

A k -additive game v^* with Möbius transform m^* belongs to $\mathcal{C}^k(v)$ if and only if it satisfies the system

$$\sum_{\substack{K \subseteq A \\ |K| \leq k}} m^*(K) \geq \sum_{K \subseteq A} m(K), \quad A \in 2^N \setminus \{\emptyset, N\} \quad (2)$$

$$\sum_{\substack{K \subseteq N \\ |K| \leq k}} m^*(K) = v(N). \quad (3)$$

271 The number of variables is $N(k) := \binom{n}{1} + \dots + \binom{n}{k}$, but due to (3), this
 272 gives rise to a $(N(k) - 1)$ -dim closed polyhedron. (2) is a system of $2^n - 2$
 273 inequalities. The polyhedron is convex since the convex combination of any
 274 two elements of the core is still in the core, but it is not bounded in general.
 275 To see this, consider the simple following example.

EXAMPLE 5: Consider $n = 3$, $k = 2$, and a game v defined by its Möbius transform m with $m(i) = 0.1$, $m(ij) = 0.2$ for all $i, j \in N$, and $m(N) = 0.1$.

Then the system of inequalities defining the 2-additive core is:

$$\begin{aligned} m^*(1) &\geq 0.1 \\ m^*(2) &\geq 0.1 \\ m^*(3) &\geq 0.1 \\ m^*(1) + m^*(2) + m^*(12) &\geq 0.4 \\ m^*(1) + m^*(3) + m^*(13) &\geq 0.4 \\ m^*(2) + m^*(3) + m^*(23) &\geq 0.4 \\ m^*(1) + m^*(2) + m^*(3) + m^*(12) + m^*(13) + m^*(23) &= 1. \end{aligned}$$

276 Let us write for convenience $m^* := (m^*(1), m^*(2), m^*(3), m^*(12), m^*(13), m^*(23))$.

277 Clearly $m_0^* := (0.2, 0.1, 0.1, 0.2, 0.2, 0.2)$ is a solution, as well as
 278 $m_0^* + t(1, 0, 0, -1, 0, 0)$ for any $t \geq 0$. Hence $(1, 0, 0, -1, 0, 0)$ is
 279 a ray, and the core is unbounded.

For the monotone core, from Prop. 1 (i) there is an additional system of $n2^{n-1}$ inequalities

$$\sum_{\substack{K \in [i, L] \\ |K| \leq k}} m^*(K) \geq 0, \quad \forall i \in N, \forall L \ni i. \quad (4)$$

For monotone games, Miranda and Grabisch [14] have proved that the Möbius transform is bounded as follows:

$$-\binom{|A| - 1}{l'_{|A|}} v(N) \leq m(A) \leq \binom{|A| - 1}{l_{|A|}} v(N), \quad \forall A \subseteq N,$$

where $l_{|A|}, l'_{|A|}$ are given by:

- (i) $l_{|A|} = \frac{|A|}{2}$, and $l'_{|A|} = \frac{|A|}{2} - 1$ if $|A| \equiv 0 \pmod{4}$
- (ii) $l_{|A|} = \frac{|A| - 1}{2}$, and $l'_{|A|} = \frac{|A| - 3}{2}$ or $l'_{|A|} = \frac{|A| + 1}{2}$ if $|A| \equiv 1 \pmod{4}$
- (iii) $l_{|A|} = \frac{|A|}{2} - 1$, and $l'_{|A|} = \frac{|A|}{2}$ if $|A| \equiv 2 \pmod{4}$
- (iv) $l_{|A|} = \frac{|A| - 3}{2}$ or $l_{|A|} = \frac{|A| + 1}{2}$, and $l'_{|A|} = \frac{|A| - 1}{2}$ if $|A| \equiv 3 \pmod{4}$.

Since $v(N)$ is fixed and bounded, the monotone k -additive core is always bounded.

For $\mathcal{C}_{\infty}^k(v)$, using Prop. 1 (ii) system (4) is replaced by a system of $N(k) - n$ inequalities:

$$m^*(K) \geq 0, \quad K \in \mathcal{P}_*^k(N), |K| > 1. \quad (5)$$

Since in addition we have $m^*(\{i\}) \geq m(\{i\})$, $i \in N$ coming from (2), m^* is bounded from below. Then (3) forces m^* to be bounded from above, so that $\mathcal{C}_{\infty}^k(v)$ is bounded.

In summary, we have the following.

Proposition 11 *For any game v , $\mathcal{C}^k(v)$, $\mathcal{MC}^k(v)$ and $\mathcal{C}_{\infty}^k(v)$ are closed convex $(N(k)-1)$ -dimensional polyhedra. Only $\mathcal{MC}^k(v)$ and $\mathcal{C}_{\infty}^k(v)$ are always bounded.*

The following result about rays of $\mathcal{C}^k(v)$ is worthwhile to be noted.

294 **Proposition 12** *The components of rays of $\mathcal{C}^k(v)$ do not depend on v , but*
 295 *only on k and n .*

296 **Proof:** For any polyhedron defined by a system of m inequalities and n
 297 variables (including slack variables) $\mathbf{Ax} = \mathbf{b}$, it is well known that its conical
 298 part is given by $\mathbf{Ax} = \mathbf{0}$, and that rays (also called basic feasible directions)
 299 are particular solutions of the latter system with $n - m$ non basic components
 300 all equal to zero but one (see, e.g., [6]). Hence, components of rays do not
 301 depend on $\mathbf{b}$.

302 Applied to our case, this means that components of rays do not depend on v ,
 303 but only on k and n . ■

304

305 4.2 A Shapley-Ichiishi-like result

We turn now to the characterization of vertices induced by achievable families.
 Let v be a game on N , m its Möbius transform, and $\prec$ be a total order on
 $\mathcal{P}_*^k(N)$. We define a k -additive game $v_\prec$ by its Möbius transform as follows:

$$m_\prec(B) := \begin{cases} \sum_{A \in \mathcal{A}(B)} m(A), & \text{if } \mathcal{A}(B) \neq \emptyset \\ 0, & \text{else} \end{cases} \quad (6)$$

306 for all $B \in \mathcal{P}_*^k(N)$, and $m_\prec(B) := 0$ if $B \notin \mathcal{P}_*^k(N)$.

307 Due to Prop. 3, $m_\prec$ satisfies $\sum_{B \subseteq N} m_\prec(B) = \sum_{B \subseteq N} m(B) = v(N)$, hence
 308 $v_\prec(N) = v(N)$.

309 This definition is a generalization of the definition of ϕ^σ or ϕ^C (see Sec. 2).
 310 Indeed, denoting by σ the permutation on N corresponding to $\prec$, we get:

$$\begin{aligned}
m_{\prec}(\{\sigma(i)\}) &= \sum_{A \subseteq \{\sigma(1), \dots, \sigma(i-1)\}} m(A \cup \sigma(i)) \\
&= \sum_{A \subseteq \{\sigma(1), \dots, \sigma(i)\}} m(A) - \sum_{A \subseteq \{\sigma(1), \dots, \sigma(i-1)\}} m(A) \\
&= v(\{\sigma(1), \dots, \sigma(i)\}) - v(\{\sigma(1), \dots, \sigma(i-1)\}) = \phi^{\sigma}(\{\sigma(i)\}) = m^{\sigma}(\{\sigma(i)\}),
\end{aligned}$$

311 where m^{σ} is the Möbius transform of ϕ^{σ} (see Sec. 2).

312 **Proposition 13** *Assume that $\mathcal{A}(B)$ is a nonempty lattice. Then $v_{\prec}(\check{B}) =$*
313 *$v(\check{B})$ if and only if $\{\mathcal{A}(C) \mid C \in \mathcal{P}_{*}^k(N), C \subseteq \check{B}, \mathcal{A}(C) \neq \emptyset\}$ is a partition of*
314 *$\mathcal{P}(\check{B}) \setminus \{\emptyset\}$.*

Proof: We have by Eq. (6)

$$v_{\prec}(\check{B}) = \sum_{\substack{C \subseteq \check{B} \\ C \in \mathcal{P}_{*}^k(N) \\ \mathcal{A}(C) \neq \emptyset}} m_{\prec}(C) = \sum_{\substack{C \subseteq \check{B} \\ C \in \mathcal{P}_{*}^k(N) \\ \mathcal{A}(C) \neq \emptyset}} \sum_{K \in \mathcal{A}(C)} m(K). \quad (7)$$

315 On the other hand, $v(\check{B}) = \sum_{K \subseteq \check{B}} m(K)$. To ensure $v_{\prec}(\check{B}) = v(\check{B})$ for any
316 v , every $K \subseteq \check{B}$ must appear exactly once in the last sum of (7), which is
317 equivalent to the desired condition. ■

318

319 The following is immediate from Prop. 13 and 7.

320 **Corollary 3** *Assume $\prec$ is compatible, and consider a nonempty achievable*
321 *family $\mathcal{A}(B)$. Then $v_{\prec}(\check{B}) = v(\check{B})$.*

322 **Proposition 14** *Let us suppose that all nonempty achievable families are lat-*
323 *tices. Then v k -monotone implies that $v_{\prec}$ is infinitely monotone.*

Proof: It remains to show that $m_{\prec}(B) \geq 0$ for any B such that $1 < |B| \leq k$.

For all such B satisfying $\mathcal{A}(B) \neq \emptyset$,

$$m_{\prec}(B) = \sum_{A \in \mathcal{A}(B)} m(A) = \sum_{A \in [B, \check{B}]} m(A).$$

324 Since $1 < |B| \leq k$, by Prop. 1, it follows from k -monotonicity that $m_{\prec}(B) \geq 0$
 325 for all $B \in \mathcal{P}_*^k(N)$. ■

326

327 The next corollary follows from Prop. 6.

328 **Corollary 4** *Let us suppose that $\prec$ is compatible. Then v k -monotone implies*
 329 *that $v_{\prec}$ is infinitely monotone.*

330 **Theorem 2** *v is $(k+1)$ -monotone if and only if for all compatible orders $\prec$,*
 331 *$v_{\prec}(A) \geq v(A)$, $\forall A \subseteq N$.*

Proof: For any compatible order $\prec$, and any $A \subseteq N$, $A \neq \emptyset$, by compatibility and Prop. 6, we can write

$$v_{\prec}(A) = \sum_{\substack{B \subseteq A \\ B \in \mathcal{P}_*^k(N) \\ \mathcal{A}(B) \neq \emptyset}} \sum_{C \in [B, \check{B}]} m(C). \quad (8)$$

Let $C \subseteq A$. Then by Prop. 3, $C \in \mathcal{A}(B)$ for some $B \subseteq A$. Indeed $B \subseteq C \subseteq A$. Hence (8) writes

$$v_{\prec}(A) = v(A) + \sum_{\substack{B \subseteq A \\ B \in \mathcal{P}_*^k(N) \\ \mathcal{A}(B) \neq \emptyset}} \sum_{\substack{C \in [B, \check{B}] \\ C \not\subseteq A}} m(C). \quad (9)$$

($\Rightarrow$) Let us take any compatible order $\prec$. By (9), it suffices to show that

$$\sum_{\substack{C \in [B, \check{B}] \\ C \not\subseteq A}} m(C) \geq 0, \quad \forall B \subseteq A, B \in \mathcal{P}_*^k(N), \mathcal{A}(B) \neq \emptyset. \quad (10)$$

332 For simplicity define $\mathcal{C}$ as the set of subsets C satisfying the condition in the
 333 summation in (10). If $\check{B} \subseteq A$, then $\mathcal{C} = \emptyset$, and so (10) holds for such B 's.
 334 Assume then that $\check{B} \setminus A \neq \emptyset$. Let us take $i \in \check{B} \setminus A$. Then $C_0 := B \cup i$ is
 335 a minimal element of $\mathcal{C}$, of cardinality $1 < |B| + 1 \leq k + 1$. Observe that

336 $[C_0, \check{B}] \subseteq \mathcal{C}$, and that it is a Boolean sublattice of $[B, \check{B}]$. Hence, $(k+1)$ -
 337 monotonicity implies that $\sum_{C \in [C_0, \check{B}]} m(C) \geq 0$ (see Prop. 1).

338 Consider $j \in \check{B} \setminus A$, $j \neq i$. If no such j exists, then $[C_0, \check{B}] = \mathcal{C}$, and we
 339 have shown (10) for such B 's. Otherwise, define $C_1 := B \cup j$ and the interval
 340 $[C_1, \check{B} \setminus i]$, which is disjoint from $[C_0, \check{B}]$. Applying again $(k+1)$ -monotonicity
 341 we deduce that $\sum_{D \in [C_1, \check{B}]} m(D) \geq 0$. Continuing this process until all elements
 342 of $\check{B} \setminus A$ have been taken, the set $\mathcal{C}$ has been partitioned into intervals $[B \cup$
 343 $i, \check{B}]$, $[B \cup j, \check{B} \setminus i]$, $[B \cup k, \check{B} \setminus \{i, j\}]$, $\dots$, $[B \cup l, A \cup l]$ where the sum of $m(C)$
 344 over these intervals is non negative by $(k+1)$ -monotonicity. Hence (10) holds
 345 in any case and the sufficiency is proved.

346 ($\Leftarrow$) Consider $K, L \subseteq N$ such that $1 < |K| \leq k+1$ and $L \supseteq K$. We have to
 347 prove that $\sum_{C \in [K, L]} m(C) \geq 0$. Without loss of generality, let us assume for
 348 simplicity that $K := \{i, i+1, \dots, l\}$ and $L := \{1, \dots, l\}$, with $l-k \leq i < l \leq n$.
 349 Define $B := K \setminus i = \{i+1, \dots, l\}$ and $A := L \setminus i$. Take a total order on $\mathcal{P}_*^k(N)$
 350 as follows:

- 351 (i) put first all subsets in $\mathcal{P}_*^k(L)$, with increasing cardinality, except B which
 352 is put the last
- 353 (ii) then put remaining subsets in $\mathcal{P}_*^k(N)$ such that they form a compatible
 354 order (for example: consider the above fixed sequence in $\mathcal{P}_*^k(L)$ augmented
 355 with the empty set as first element of the sequence, then take any subset
 356 D in $N \setminus L$ belonging to $\mathcal{P}_*^k(N)$, and add it to any subset of the sequence,
 357 discarding subsets not in $\mathcal{P}_*^k(N)$. Do this for any subset D of $N \setminus L$).
- 358 (iii) subsets in $\mathcal{P}_*^k(L)$ with same cardinality are ordered according to the lex-
 359 icographic order, which means in particular $1 \prec 2 \prec \dots \prec l$.

360 One can check that such an order is compatible¹. By construction, we have

¹ For example, with $n = 5, l = 4, i = 3, k = 3$:

$$1 \prec 2 \prec 3 \prec 12 \prec 13 \prec 14 \prec 23 \prec 24 \prec 34 \prec 123 \prec 124 \prec 134 \prec 234 \prec 4 \prec 5 \prec 51 \prec 52 \dots$$

$\mathcal{A}(B) = [B, L]$. Indeed, for any $C \in \mathcal{A}(B)$, any subset of C in $\mathcal{P}_*^k(N)$ is ranked
 before B . Moreover, $[K, L] = [B \cup i, L] = \{C \in \mathcal{A}(B) \mid C \not\subseteq A\}$. Now, take
 any $B' \neq B$ in $\mathcal{P}_*^k(L)$ such that $B' \subseteq A$. Let us prove that any $C \in \mathcal{A}(B')$
 is such that $i \notin C$, or equivalently $C \subseteq A$. Indeed, up to the fact that B is
 ranked last, the sequence $\mathcal{P}_*^k(L)$ forms a strongly compatible order. Adapting
 slightly Prop. 9, it is easy to see that if $|B'| < k$, then either $\check{B}' = B'$ or
 $\mathcal{A}(B') = \emptyset$, the latter arising if $B' \supset B$. Then trivially any $C \in \mathcal{A}(B')$ satisfies
 $C \subseteq A$. Assume now $|B'| = k$. If B' contains some $j \prec i$, then $B' \cup i$ cannot
 belong to $\mathcal{A}(B')$ since by lexicographic ordering $B' \cup i \setminus j$ is ranked after B' ,
 which implies that for any $C \in \mathcal{A}(B')$, $i \notin C$. Hence, the condition $i \in C$ can
 be true for some $C \in \mathcal{A}(B')$ only if all elements of B' are ranked after i . But
 since $B = \{i + 1, \dots, l\}$, this implies that either $B' = B$, a contradiction, or
 B' does not exist (if $|B| < k$).

Let us apply the dominance condition for $v_{\prec}(A)$. Using (9), dominance is equivalent to write:

$$\sum_{\substack{B \subseteq A \\ B \in \mathcal{P}_*^k(N) \\ \mathcal{A}(B) \neq \emptyset}} \sum_{\substack{C \in [B, \check{B}] \\ C \not\subseteq A}} m(C) \geq 0.$$

Using the above, this sum reduces to $\sum_{C \in [K, L]} m(C) \geq 0$. This finishes the proof. ■

The following is an interesting property of the system $\{(2), (3)\}$.

Proposition 15 *Let $\prec$ be a compatible order. Then the linear system of equalities $v_{\prec}(\check{B}) = v(\check{B})$, for all $\check{B}$'s induced by $\prec$, is triangular with no zero on the diagonal, and hence has a unique solution.*

Proof: We consider w.l.o.g. that $1 \prec 2 \prec \dots \prec n$ and consider the binary order $\prec^2$ for ordering variables $m^*(B)$.

383 Delete all variables such that $\mathcal{A}(B) = \emptyset$, and consider the list of subsets
 384 in $\mathcal{P}_*^k(N)$ corresponding to non deleted variables. Take all B 's in the list,
 385 and their corresponding $\check{B}$'s (always exist since by compatibility, $\mathcal{A}(B)$ is a
 386 lattice). They are all different by Prop. 3, so we get a linear system of the
 387 same number of equations (namely $v_{\prec}(\check{B}) = v(\check{B})$) and variables. Take one
 388 particular equation corresponding to B . Then variables used in this equation
 389 are necessarily $m^*(B)$ itself (because $\check{B} \supseteq B$), and some variables ranked
 390 before B in the binary order. Indeed, if $\check{B} = B$, then all variables used in the
 391 equation are ranked before B by $\prec^2$. If $\check{B} \neq B$, supersets B' of B in $\mathcal{P}_*^k(N)$
 392 are ranked after B by $\prec^2$ (because $\prec^2$ is $\subseteq$ -compatible), and ranked before
 393 B by $\prec$ (otherwise $\mathcal{A}(B)$ would not contain $\check{B}$), but since they contain B ,
 394 necessarily $\mathcal{A}(B') = \emptyset$, so corresponding variables are deleted.

395 Hence the system is triangular. ■

396

397 Note that the proof holds under the condition that all achievable families are
 398 lattices, so compatibility is even not necessary.

399 **Theorem 3** *Let v be a $(k+1)$ -monotone game. Then*

- 400 (i) *If $\prec$ is strongly compatible, then $v_{\prec}$ is a vertex of $\mathcal{C}^k(v)$.*
- 401 (ii) *If $\prec$ is compatible, then $v_{\prec}$ is a vertex of $\mathcal{C}_{\infty}^k(v)$.*

402 **Proof:** By standard results on polyhedra, it suffices to show that $v_{\prec}$ is an
 403 element of $\mathcal{C}^k(v)$ (resp. $\mathcal{C}_{\infty}^k(v)$) satisfying at least $N(k)-1$ linearly independent
 404 equalities among (2) (resp. among (2) and (5)). Assume $\prec$ is compatible. Then
 405 by Cor. 4, $v_{\prec}$ is infinitely monotone, and it dominates v by $(k+1)$ -monotonicity
 406 (Th. 2). Moreover, for any $B \in \mathcal{P}_*^k(N)$, $\mathcal{A}(B)$ is either empty or a lattice, hence
 407 either $m_{\prec}(B) = 0$ or $v_{\prec}(\check{B}) = v(\check{B})$ by Cor. 3. Since if $|B| = 1$, $\mathcal{A}(B) \neq \emptyset$,
 408 this gives $N(k)$ equalities in the system defining $\mathcal{C}_{\infty}^k(v)$, including (3), hence

we have the exact number of equalities required, which form a nonsingular system by Prop 15, and (ii) is proved. If the order is strongly compatible, then all achievable families are lattices, which proves the result for $\mathcal{C}^k(v)$, since again by Prop. 15, the system is nonsingular. ■

413

REMARK 1: Vertices induced by (strongly) compatible orders are also vertices of the monotone k -additive core. They are induced only by dominance constraints, not by monotonicity constraints.

REMARK 2: Cor. 3 generalizes Prop. 2, while Theorems 2 and 3 generalize the Shapley-Ichiishi results summarized in Th. 1. Indeed, recall that convexity is 2-monotonicity. Then clearly Th. 2 is a generalization of (i) $\Rightarrow$ (ii) of Th. 1, and Th. 3 (i) is a part of (iv) in Th. 1. But as it will become clear below, all vertices are not recovered by achievable families, mainly because they can induce only infinitely monotone games. In particular, $\mathcal{MC}^k(v)$ contains many more vertices.

Let us examine more precisely the number of vertices induced by strongly compatible orders. In fact, there are much fewer than expected, since many strongly compatible orders lead to the same $v_{\prec}$. The following is a consequence of Prop. 10.

Corollary 5 *The number of vertices of $\mathcal{C}^k(v)$ given by strongly compatible orders is at most $\frac{n!}{k!}$.*

Proof: Given the order $1 \prec 2 \prec \dots \prec n$, a permutation over the last k singletons would not change the collection $\check{\mathcal{B}}$. ■

432

Note that when $k = 1$, we recover the fact that vertices are induced by all permutations, and that with $k = n$, we find only one vertex (which is in fact

the only vertex of $\mathcal{C}^n(v)$, which is v itself (use Prop. 10 and the definition of $m_{\prec}$).

4.3 Other vertices

In this last section we give some insights about other vertices. Even for the (non monotonic) k -additive core, in general for $k \neq 1, n$, not all vertices are induced by strongly compatible orders. However, for the case $k = n - 1$, it is possible to find all vertices of $\mathcal{C}^k(v)$. For $1 < k < n - 1$ and also for the monotonic core, the problem becomes highly combinatorial.

Theorem 4 *Let v be any game in $\mathcal{G}(N)$, with Möbius transform m .*

(i) *If $m(N) > 0$, $\mathcal{C}^{n-1}(v)$ contains exactly 2^{n-1} (if n is even) or $2^{n-1} - 1$ (if n is odd) vertices, among which n vertices come from strongly compatible orders. They are given by their Möbius transform:*

$$m_{B_0}^*(K) = \begin{cases} m(K), & \text{if } K \not\supseteq B_0 \\ m(K) + (-1)^{|K \setminus B_0|} m(N), & \text{else} \end{cases}$$

for all $B_0 \subset N$ such that $|N \setminus B_0|$ is odd.

(ii) *If $m(N) = 0$, then there is only one vertex, which is v itself.*

(iii) *If $m(N) < 0$, $\mathcal{C}^{n-1}(v)$ contains exactly $2^{n-1} - 1$ (if n is odd) or $2^{n-1} - 2$ (if n is even) vertices, of which none comes from a strongly compatible order. They are given by their Möbius transform:*

$$m_{B_0}^*(K) = \begin{cases} m(K), & \text{if } K \not\supseteq B_0 \\ m(K) - (-1)^{|K \setminus B_0|} m(N), & \text{else} \end{cases}$$

for all $B_0 \subset N$ such that $|N \setminus B_0|$ is even.

Proof: We assume $m(N) \geq 0$ (the proof is much the same for the case $m(N) \leq 0$). We consider the system of $2^n - 1$ inequalities $\{(2), (3)\}$, which has

$N(n-1) = 2^n - 2$ variables. We have to fix $2^n - 2$ equalities, among which (3), so we have to choose only one inequality in (2) to remain strict, say for $B_0 \subset N$, $B_0 \neq \emptyset$:

$$\sum_{K \subseteq B_0} m^*(K) > \sum_{K \subseteq B_0} m(K). \quad (11)$$

From the definition of the Möbius transform, we have

$$0 = m^*(N) = \sum_{K \subseteq N} (-1)^{|N \setminus K|} v^*(K).$$

447 Note that for any $\emptyset \neq K \subseteq N$, $v^*(K)$ is the left member of some inequality
 448 or equality of the system. Hence, by turning all inequalities into equalities,
 449 we get, by doing the above summation on the system, $0 = m(N)$. Hence, if
 450 $m(N) = 0$, there is only one vertex, which is v itself, otherwise taking equality
 451 everywhere gives a system with no solution. Since strict inequality holds only
 452 for $B_0 \subset N$, we get instead $0 > m(N)$ if $|N \setminus B_0|$ is even, and $0 < m(N)$ if
 453 $|N \setminus B_0|$ is odd. The first case is impossible by assumption on $m(N)$, so only
 454 the case where $|N \setminus B_0|$ odd can produce a vertex. Note that if $|B_0| = n - 1$,
 455 we recover all n vertices induced by strongly compatible orders. In total we
 456 get $\binom{n}{n-1} + \binom{n}{n-3} + \dots + \binom{n}{1} = 2^{n-1}$ potential different vertices when n is even,
 457 and $2^{n-1} - 1$ when n is odd. Clearly, there is no other possibility.

It remains to show that the corresponding system of equalities is non singular, and eventually to solve it. Assume $B_0 \subset N$ in (11) is chosen. From the linear system of equalities we easily deduce $m^*(K) = m(K)$ for all $K \not\supseteq B_0$. Substituting into all equations, the system reduces to

$$\begin{aligned} \sum_{K \subseteq B \setminus B_0} m^*(B_0 \cup K) &= \sum_{K \subseteq B \setminus B_0} m(B_0 \cup K), \quad \forall B \supset B_0, B \neq N \\ \sum_{K \subset N \setminus B_0} m^*(B_0 \cup K) &= \sum_{K \subset N \setminus B_0} m(B_0 \cup K) + m(N). \end{aligned}$$

B_0 being present everywhere, we may rename all variables after deleting B_0 , i.e., we set $N' = N \setminus B_0$, $m'(A) := m(A \cup B_0)$ and $m'^*(A) = m^*(A \cup B_0)$, for

all $A \subseteq N'$. The system becomes

$$\begin{aligned}\sum_{K \subseteq B} m'^*(K) &= \sum_{K \subseteq B} m'(K), \quad \forall B \subset N' \\ \sum_{K \subset N'} m'^*(K) &= \sum_{K \subset N'} m'(K) + m'(N').\end{aligned}$$

458 Summing equations of the system as above, i.e., computing $\sum_{B \subseteq N'} (-1)^{|N' \setminus B|} \sum_{K \subseteq B} m'^*(K)$,
 459 we get $m'^*(\emptyset) = m'(\emptyset) + m'(N')$, or equivalently $m^*(B_0) = m(B_0) + m(N)$.
 460 Substituting in the above system, we get a system which is triangular (use, e.g.,
 461 Prop. 15 with $k = n = n'$). We get easily $m^*(K) = m(K) + (-1)^{|K \setminus B_0|} m(N)$,
 462 for all $K \supseteq B_0$. ■

463

464 References

- 465 [1] E. Algaba, J. M. Bilbao, R. van den Brink, and A. Jiménez-Losada. Cooperative
 466 games on antimatroids. *Discrete Mathematics*, 282:1–15, 2004.
- 467 [2] J.-P. Barthélemy. Monotone functions on finite lattices: an ordinal approach to
 468 capacities, belief and necessity functions. In J. Fodor, B. De Baets, and P. Perny,
 469 editors, *Preferences and Decisions under Incomplete Knowledge*, pages 195–208.
 470 Physica Verlag, 2000.
- 471 [3] J. M. Bilbao, N. Jiménez, E. Lebrón, and H. Peters. The selectope for games
 472 with partial cooperation. *Discrete Mathematics*, 216:11–27, 2000.
- 473 [4] O. Bondareva. Some applications of linear programming to the theory of
 474 cooperative games. *Problemy Kibernet*, 10:119–139, 1963.
- 475 [5] A. Chateauneuf and J.-Y. Jaffray. Some characterizations of lower probabilities
 476 and other monotone capacities through the use of Möbius inversion.
 477 *Mathematical Social Sciences*, 17:263–283, 1989.
- 478 [6] V. Chvátal. *Linear Programming*. Freeman and Company, 1983.

- 479 [7] U. Faigle and W. Kern. On the core of ordered submodular cost games.
480 *Mathematical Programming*, 87:483–499, 2000.
- 481 [8] U. Faigle, W. Kern, S. Fekete, and W. Hochstättler. The nucleon of cooperative
482 games and an algorithm for matching games. *Mathematical Programming*,
483 83:195–211, 1998.
- 484 [9] S. Fujishige. *Submodular functions and optimization*. Elsevier, 2nd edition
485 edition, 2005.
- 486 [10] M. Grabisch. k -order additive discrete fuzzy measures and their representation.
487 *Fuzzy Sets and Systems*, 92:167–189, 1997.
- 488 [11] J. C. Harsanyi. A simplified bargaining model for the n -person cooperative
489 game. *International Economic Review*, 4:194–220, 1963.
- 490 [12] T. Ichiishi. Super-modularity: applications to convex games and to the greedy
491 algorithm for LP. *J. Econom. Theory*, 25:283–286, 1981.
- 492 [13] L. Lovász. Submodular function and convexity. In A. Bachem, M. Grötschel,
493 and B. Korte, editors, *Mathematical programming. The state of the art*, pages
494 235–257. Springer Verlag, 1983.
- 495 [14] P. Miranda and M. Grabisch. Optimization issues for fuzzy measures. *Int. J.*
496 *of Uncertainty, Fuzziness, and Knowledge-Based Systems*, 7(6):545–560, 1999.
- 497 [15] P. Miranda and M. Grabisch. k -balanced capacities. In *Proc. of the Int. Fuzzy*
498 *Systems Association World Congress (IFSA)*, pages 455–460, Beijing, China,
499 July 2005.
- 500 [16] G. C. Rota. On the foundations of combinatorial theory I. Theory of Möbius
501 functions. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*,
502 2:340–368, 1964.
- 503 [17] L. S. Shapley. Core of convex games. *Int. J. Game Theory*, 1:11–26, 1971.
- 504 [18] J. Von Neuman and O. Morgenstern. *Theory of Games and Economic Behavior*.
505 Princeton University Press, 1944.

506 [19] P. Walley. *Statistical Reasoning with Imprecise Probabilities*. Chapman and
507 Hall, London, 1991.